




SUPPLEMENTARY FIGURES




Supplementary Figure 1. The results of a sensitivity analysis for 90-day favorable functional outcomes in acute ischemic stroke patients who underwent endovascular therapy.




Supplementary Figure 2. The results of a sensitivity analysis for recanalization rates in acute ischemic stroke patients who underwent endovascular therapy.




Supplementary Figure 3. The results of a sensitivity analysis for sICH rates in acute ischemic stroke patients who underwent endovascular therapy.




Supplementary Figure 4. The results of a sensitivity analysis for 90-day mortality in acute ischemic stroke patients who underwent endovascular therapy.




Supplementary Figure 5. A funnel plot for assessing publication bias in regards to studies reporting 90-day favorable functional outcomes in acute ischemic stroke patients who underwent endovascular therapy.



Supplementary Figure 6. A funnel plot for assessing publication bias in regards to studies reporting recanalization rates in acute ischemic stroke patients who underwent endovascular therapy.



Supplementary Figure 7. A funnel plot for assessing publication bias in regards to studies reporting 90-day mortality in acute ischemic stroke patients who underwent endovascular therapy.



Supplementary Figure 8. A funnel plot for assessing publication bias in regards to studies reporting sICH in acute ischemic stroke patients who underwent endovascular therapy.